

The morphology-density relation in X-ray bright galaxy groups

Stephen F. Helsdon^{1,2*} and Trevor J. Ponman²

¹ *The Observatories of the Carnegie Institute of Washington, 813 Santa Barbara Street, Pasadena, CA 91101*

² *School of Physics and Astronomy, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK*

Accepted 2002 ??; Received 2001 ??; in original form 2001 ??

ABSTRACT

We have examined the morphological make-up of X-ray bright groups. The brighter galaxies in these groups exhibit clear morphology-density and morphology-radius relations. The group morphology-density relation is offset from the cluster relation in the sense that at a given surface density, X-ray bright groups have a lower spiral fraction. After correcting for projection effects the morphology-3D density relation is still shifted towards fewer spirals for a given 3D density, in comparison with clusters. A simple model which corrects the group data for the effects of projection and for the expected higher merging rate in groups, brings the morphology-density relation into good agreement with that of clusters, suggesting that the relation may be driven by two-body interactions. The fraction of S0 galaxies in these X-ray bright groups is at least as high as that observed in nearby clusters. Given the low velocity dispersion of groups, this indicates that ram pressure stripping is not the dominant mechanism for S0 formation.

Key words: X-rays: galaxies: clusters – X-rays: galaxies – intergalactic medium – galaxies: clusters: general – galaxies: evolution

1 INTRODUCTION

The well known relationship between galaxy morphology and density (e.g. Dressler 1980; Dressler et al. 1997) suggests that the environment of a galaxy may have a significant effect on its evolution. By examining the properties of galaxies as a function of environment it should be possible to gain important insights into the evolution of both galaxies and galaxy systems. Particularly interesting is the group environment, which is typically made up of between three and a few tens of galaxies. These systems are gravitationally bound systems in which the density and velocities of the member galaxies suggest that mergers and interactions are more common than in clusters or in the field (e.g. Mamon 1992, 2000a). In addition, the majority of galaxies are found in a group environment (Tully 1987), and many galaxies in clusters will once have been part of groups.

Perhaps the most striking connection between group and galaxy properties is the fact that almost all X-ray bright groups contain a bright early-type galaxy located at the spatial and kinematic centre of the group (Zabludoff & Mulchaey 1998). This central galaxy is presumably the re-

sult of early merging activity in the collapsing group core (Governato et al. 1996) and given the high expected rates of mergers in present day groups, this central galaxy may still be rapidly growing through mergers and accretion. Other galaxies in the group are also likely to be affected by the group environment. Postman & Geller (1984) and Ramella et al. (1999) have shown that galaxies in groups follow a morphology-density relation. However, it is not clear if this is the same relation as for clusters as some X-ray bright groups appear to have spiral fractions consistent with those found in rich clusters (Zabludoff & Mulchaey 1998).

It is also known that there is a connection between the morphological make-up of groups and the presence of diffuse X-ray emission (e.g. Pildis et al. 1995; Henry et al. 1995; Mulchaey et al. 1996), in the sense that X-ray detected groups tend to be spiral poor, whilst undetected groups contain more late-types. In this and a companion paper (Helsdon & Ponman 2002) we aim to look in some detail at the relationship between the X-ray and optical properties of X-ray bright galaxy groups. Here we focus on the morphological makeup of these systems, and examine the morphology-density relation of X-ray bright groups.

* E-mail: sfh@ociw.edu




Figure 1. Spiral fraction (including irregulars) versus radius (in units of the virial radius, R_V) for X-ray bright groups. Each radial bin contains at least 75 galaxies.

2 THE SAMPLE

The group sample used here is the sample of 24 X-ray bright groups originally studied by (Helsdon & Ponman 2000a). The X-ray properties of this sample are discussed extensively in Helsdon & Ponman (2000a) and Helsdon & Ponman (2000b), but the important thing for the purposes of this work is that all these groups have detected, extended diffuse emission, associated with the group. The presence of a hot intragroup medium indicates that the group is a real gravitationally bound object, rather than just a chance superposition of a few galaxies — potentially a serious problem with optically selected groups.

We also need to determine the galaxy memberships of these groups. Unfortunately, it is not satisfactory to simply use the galaxy memberships as given in the original group catalogue, since the galaxies were originally selected from different sources which in general have different selection criteria. This problem is addressed in Helsdon & Ponman (2002), and here we use the same technique. This procedure is summarised below — for full details see Helsdon & Ponman (2002). For each group we search the NASA/IPAC Extragalactic Database (NED) and the Lyon-Meudon Extragalactic Database (LEDa) for galaxies lying within the group virial radius in projection on the sky, and having recession velocities within three times the group velocity dispersion from the catalogued group mean. The centre position used for each group is the centre of the X-ray emission. In most cases this position is very close to the position of the central galaxy, with the three exceptions being bimodal systems in which the X-ray centre falls roughly between the two main galaxies. It is possible that the centres used in the bimodal groups could act to obscure radial trends present in the other groups. However, as we average over all groups when looking at radial trends, and the bimodal systems only make up a small fraction of the sample, this is unlikely to have a significant effect on any results.

To ensure an even luminosity cut in all groups we use

the galaxy luminosity function of Zabludoff & Mulchaey (2000) (for X-ray bright groups) to employ two magnitude cuts — one which should include 50% of the total optical group light and one which should include 90% of the total group light. For the 50% cut Helsdon & Ponman (2002) show that for the typical group in this sample the membership should be almost 100% complete, whilst approximately 35% of the total group light is missing from the 90% cut sample. Morphological types are taken firstly from NED if available, and then LEDa.

It should be noted that the group sample used here should not be regarded as being statistically complete in any way. However, we do not believe that this will introduce any particular bias, other than the fact that since we only use groups with detected diffuse X-ray emission, we do not include systems with undetectably faint intergalactic gas. The group sample should rather be regarded as a reasonably representative sample of X-ray bright groups.

3 THE MORPHOLOGY-DENSITY RELATION

The overall spiral fraction of all these X-ray bright groups combined is 0.425 ± 0.035 , however this spiral fraction drops to 0.30 ± 0.05 if the sample is restricted to galaxies brighter than the 50% luminosity cut. Thus these groups have lower spiral fractions than found in the field (~ 0.6 — Postman & Geller 1984; Whitmore et al. 1993), particularly amongst the more luminous galaxies.

The spiral fraction also appears to increase with radius in these systems. In Figure 1 we plot the spiral fraction calculated from all galaxies in all groups, as a function of radius (in units of the virial radius). This plot shows that the spiral fraction drops from about 0.6 at the outer edge (consistent with the field population) to a value of about 0.2 in the innermost regions of the groups. The spiral number fraction and the spiral light fraction are similar to one another until the central bin, where the spiral light fraction drops to




Figure 2. Spiral fraction (including irregulars) as a function of surface density (averaged over radial bins). Histogram is the relation for local clusters (Dressler et al. 1997), and the open circles with error bars are the group data. Crossed and shaded circles represent group points after estimated corrections for 3d density, and merging rates respectively (for clarity the error bars are omitted from these points).

less than half the number fraction, due to the presence of a central bright early type in almost all these groups. The spiral fraction profile for the 50% sample is similar to the 90% sample shown, but has a slightly lower spiral fraction at each point.

This morphology-radius trend is similar to that observed in galaxy clusters (Whitmore et al. 1993) and also reported in X-ray bright groups (Tran et al. 2001). We have also looked for evidence of a morphology-density relation. We cannot calculate a local surface density for each galaxy in our sample as has been done in previous cluster work (e.g. Dressler 1980; Dressler et al. 1997) as we do not have sufficient numbers of galaxies in each group. Instead we calculate the average density of each of the 4 radial bins plotted on Figure 1. We plot these points in Figure 2 along with data from the local cluster sample of Dressler et al. (1997). Note that the 50% group sample data is used, as the magnitude cut for this sample is closest to the average of that used by Dressler et al. for the local cluster data (the local cluster data is derived from an apparent magnitude cut of $V = 16.5$, which in turn corresponds to an absolute magnitude of $M_V = -20.5$ at the average cluster distance. Our 50% group sample B-band cut falls within the range of magnitude cuts for the local cluster sample and is between 0.5 and 0.9 magnitudes brighter than the average, depending on galaxy type).

As can be seen in Figure 2, the group points (circles) do indeed appear to follow a morphology-density relation. However, it is clear that the group relation is offset from the cluster data of Dressler et al.. One explanation for this offset could be that the relation is driven by 3D density, rather than by *projected* density. In a group, fewer galaxies will be projected along the line of sight onto the centre of the group. Thus if a group and cluster had the same 3D density at the centre, the projected density of the group would be lower than that of the cluster. The projected surface density (Σ) should be related to the 3D density (ρ) by, $\Sigma \propto \rho R$,

where R is the radius of the cluster. Assuming self-similarity we have, $R \propto T^{\frac{1}{2}}$, and therefore $\Sigma \propto \rho T^{\frac{1}{2}}$. So by using a mean group and mean cluster temperature, it is possible to correct for the effects of projection on the group data, and compare with the cluster data. Of the 55 clusters in the Dressler et al. local cluster sample we obtain temperatures for 31 of them from White et al. (1997) and use the mean of these as the typical cluster temperature. This gives a mean cluster temperature of $T = 3.85$ keV in comparison with a mean temperature of $T = 0.92$ keV for the group sample.

The group data corrected for the effects of 3D projection are also plotted (crossed circles) in Figure 2. It can be seen that the points have moved in such a way as to reduce the discrepancy between the group and cluster points, but there is still a significant offset between the datasets. At comparable local 3D densities, X-ray bright groups appear to have fewer spirals and more early-types than clusters. This result is inconsistent with that of Postman & Geller (1984), who found a universal morphology density relation for groups and clusters, with low density groups being more spiral rich than high density clusters, and with the higher density regions in groups having the same morphological mix as cluster regions of comparable 3D density. However, the groups used by Postman & Geller (1984) were optically selected, and it is possible that they included some spurious groups or groups at an early stage of virialisation. Both of these effects would increase the spiral fraction, and would therefore move the group data towards the cluster data. In comparison, our sample consists of collapsed, X-ray bright groups. It should also be pointed out that the trend of fewer spirals for a given 3D density is opposite to that noted by Mamon (1986) and Hickson et al. (1988) for the Hickson Compact Groups (HCG) (Hickson 1982). However, the apparent densities of HCGs are subject to complicated and subtle selection effects (e.g. Prandoni et al. 1994; Mamon 2000b) and in light of this it may not be particularly surprising that they behave in a different way.




Figure 3. a) Elliptical and b) S0 fractions as a function of surface density. The histograms are the relations for local clusters from Dressler et al. (1997), open circles are the group data and shaded circles are the group points after corrections to the surface density for the difference in merging rate between the group and cluster data.

So whilst it seems that galaxies in X-ray bright groups do follow a morphology-density relation, this relation appears to be different to that observed in clusters. However, groups are just small clusters, and it is reasonable to assume that similar processes govern the morphological makeup in both groups and clusters. The group sample here is probing the morphology-density relation in systems approximately an order of magnitude less massive than the clusters in Dressler et al. (1997), and the variations seen over this large mass range suggest that an additional parameter is needed in the morphology-density relation.

A likely candidate for this extra parameter is the merging rate – galaxies in groups have a higher merging rate than galaxies in clusters, due to their lower velocity dispersions. Using different merger cross-sections, Mamon (1992) and Makino & Hut (1997) have shown that the merging rate (k) scales as $k \propto \sigma_{cl}^{-3}$, where σ_{cl} is the cluster velocity dispersion, and Mamon (2000a) argues that this $k : \sigma_{cl}$ relation will hold for any well-behaved merger cross section. Given $\sigma_{cl} \propto T^{1/2}$, for self-similar clusters the merging rate will thus scale as $k \propto T^{-3/2}$. If we now assume that the spiral fraction in any particular environment is a function of the 3D density and the merging rate (as is expected given that the number of mergers endured by a given galaxy per unit time is $P = nk = n\langle v \Sigma(v) \rangle$, where n is the 3D number density and Σ is the velocity dependent merger cross-section, see Mamon (1992, 2000a)), we can once again scale the group densities using the mean cluster and group temperatures, in order to remove the effects of projection and merging rate. The effects of scaling the group projected densities by this factor are also shown in Figure 2. This time the group and cluster points agree fairly well, with only the densest group point departing somewhat from the cluster trend. Given the likely hierarchical development of clusters, it is important not to overinterpret this result. However, it does clearly indicate that at a given 3D density morphological transformation of galaxies is more effective within X-ray bright groups rather than within clusters.

We have also separated our group morphology-density

relation into abundances for elliptical and lenticular galaxies. These relations are compared with the equivalent cluster relations in Figure 3. Note that the group points are plotted at both the calculated surface densities and the 3D densities adjusted for their higher expected merging rate (see above). This figure clearly shows that at equivalent surface densities, the fraction of both ellipticals and lenticulars in groups tends to be higher than that in clusters. After scaling the projected densities to allow for system size and merger rate, as discussed above for spiral fractions, the points for X-ray bright groups again come into reasonable agreement with those for clusters, apart from an apparent deficit in lenticulars in groups (compared to clusters) at the highest densities.

4 DISCUSSION

Studies of the evolution of the morphology-density relation (e.g. Dressler et al. 1997) have shown that the fraction of ellipticals in clusters has not evolved significantly since $z \sim 0.5$. The elliptical fractions observed in these groups support the suggestion of Dressler et al. that the formation of ellipticals predates cluster formation, and that they are instead formed primarily in a group environment. However, the S0 fraction in clusters at $z \sim 0.5$ is about 2-3 times smaller than in present day clusters (e.g. Dressler et al. 1997; although see Andreon 1998 for a different view), suggesting that some process must have transformed the excess spirals seen in these systems into S0s by the present day.

Major mergers do not appear able to produce enough S0s in clusters (e.g. Okamoto et al. 2001a), and a number of other mechanisms have been proposed to explain the transformation of spirals to S0s: ram pressure stripping (e.g. Gunn & Gott 1972; Quilis et al. 2000), galaxy harassment (e.g. Moore et al. 1999) or unequal mass galaxy mergers (e.g. Bekki 1998). The fact that we see a comparable S0 fraction in present day X-ray bright groups and clusters, suggests that whatever process is responsible for converting spirals

to S0s in clusters also plays a significant role in creating S0s in these groups.

This would appear to rule out ram pressure stripping of the whole ISM of a galaxy as the dominant process. Indeed, since ram pressure scales as the square of the galaxy velocity within its group or cluster, ram pressure stripping should not be effective in the low velocity dispersion group environment (e.g. Zabludoff & Mulchaey 1998; Abadi et al. 1999). Furthermore, there is also evidence that ram pressure stripping cannot produce sufficient numbers of intermediate bulge-to-disk ratio galaxies in clusters (e.g. Okamoto et al. 2001b; Lanzoni 2000), where it should be more effective. However, if continuing star formation within spirals is fuelled by continuing infall of gas from a surrounding gas reservoir (still a controversial issue), then such a reservoir should be easily removed by infall into groups as well as clusters, leading to a gradual decline or ‘strangulation’ (Larson et al. 1980; Balogh et al. 2000; Balogh & Morris 2000) in star formation. There is observational evidence to suggest that this may be occurring, since late-type galaxies in groups appear to be HI-deficient (e.g. Williams & Rood 1987; Oosterloo & Iovino 1997; Verdes-Montenegro et al. 2001). The cessation of star formation would not in itself convert spirals to lenticulars, so dynamical interactions are probably involved in any case in effecting structural changes to spirals within dense environments (tidal disruption is needed to suppress spiral features and enhance the relative importance of the bulge).

It is tempting to relate the difference in behaviour in the highest density bin, between the incidence of lenticulars in X-ray bright groups and clusters, to the apparent evolution in cluster S0 content since moderate redshifts. In contrast, the elliptical content of clusters is not found to evolve, and appears to be similar to that of groups after correction for the higher merger rate in groups. It may be, for example, that galaxy harassment within the densest regions of the cluster environment has further boosted the S0 content inherited from precursor groups: galaxy harassment depends on the frequency of encounters (Moore et al. 1998), which will be higher than the merger rate, and should be greatest in the centres of clusters.

5 ACKNOWLEDGEMENTS

The authors would like to thank the referee, Gary Mamon, for his many useful comments and suggestions. This work made use of the Starlink facilities at Birmingham, the NASA/IPAC Extragalactic Database (NED — <http://nedwww.ipac.caltech.edu>) and the LEDA database (<http://leda.univ-lyon1.fr>). SFH acknowledges financial support from the University of Birmingham.

REFERENCES

Abadi M. G., Moore B., Bower R. G., 1999, MNRAS, 308, 947
 Andreon S., 1998, ApJ, 501, 533
 Balogh M. L., Morris S. L., 2000, MNRAS, 318, 703
 Balogh M. L., Navarro J. F., Morris S. L., 2000, ApJ, 540, 113

Bekki K., 1998, ApJ, 502, L133
 Dressler A., 1980, ApJ, 236, 351
 Dressler A., Oemler A., Couch W. J., Smail I., Ellis R. S., Barger A., Butcher H., Pogginnti B. M., Sharples R. M., 1997, ApJ, 490, 577
 Governato F., Tozzi P., Cavaliere A., 1996, ApJ, 458, 18
 Gunn J. E., Gott J. R., 1972, ApJ, 176, 1
 Helsdon S. F., Ponman T. J., 2000a, MNRAS, 315, 356
 Helsdon S. F., Ponman T. J., 2000b, MNRAS, 319, 933
 Helsdon S. F., Ponman T. J., 2002, MNRAS, Accepted
 Henry J., Gioia I. M., Huchra J. P., Burg R., McLean B., Böhringer H., Bower R. G., Briel U. G., Voges W., MacGillivray H., Cruddace R. G., 1995, ApJ, 449, 422
 Hickson P., 1982, ApJ, 255, 382
 Hickson P., Kindl E., Huchra J. P., 1988, ApJ, 331, 64
 Lanzoni B., 2000, PhD thesis, University of Paris 7
 Larson R. B., Tinsley B. M., Caldwell C. N., 1980, ApJ, 237, 692
 Makino J., Hut P., 1997, ApJ, 481, 83
 Mamon G. A., 1986, ApJ, 307, 426
 Mamon G. A., 1992, ApJ, 401, L3
 Mamon G. A., 2000a, in Combes F., Mamon G. A., Charmandaris V., eds, ASP Conf. Ser. 197: Dynamics of Galaxies: from the Early Universe to the Present, p. 377, astro-ph/9911333
 Mamon G. A., 2000b, in Valtonen M. J., Flynn C., eds, ASP Conf. Ser. 209: Small Galaxy Groups, p. 217, astro-ph/9909019
 Moore B., Lake G., Katz N., 1998, ApJ, 495, 139
 Moore B., Lake G., Quinn T., Stadel J., 1999, MNRAS, 304, 465
 Mulchaey J. S., Davis D. S., Mushotzky R. F., Burstein D., 1996, ApJ, 456, 80
 Okamoto T., Nagashima M., 2001a, ApJ, 547, 109
 Okamoto T., Nagashima M., 2001b, preprint, astro-ph/0108434
 Oosterloo T., Iovino A., 1997, PASA, 14, 480
 Pildis R. A., Bregman J. N., Evrard A. E., 1995, ApJ, 443, 514
 Postman M., Geller M. J., 1984, ApJ, 281, 95
 Prandoni I., Iovino A., MacGillivray H. T., 1994, AJ, 107, 1235
 Quilis V., Moore B., Bower R., 2000, Science, 288, 1617
 Ramella M., Zamorani G., Zucca E., Stirpe G. M., Vetolani G., Balkowski C., Blanchard A., Cappi A., Cayatte V., Chincarini G., Collins C., Guzzo L., Macgillivray H., Maccagni D., Maurogordato S., Merighi R., Mignoli M., Pisani A., Proust D., Scaramella R., 1999, A&A, 342, 1
 Tran K. H., Simard L., Zabludoff A. I., Mulchaey J. S., 2001, ApJ, 549, 172
 Tully R. B., 1987, ApJ, 321, 280
 Verdes-Montenegro L., Yun M. S., Williams B. A., Huchtmeier W. K., Del Olmo A., Perea J., 2001, A&A, 377, 812
 White D. A., Jones C., Forman W., 1997, MNRAS, 292, 419
 Whitmore B. C., Gilmore D. M., Jones C., 1993, ApJ, 407, 489
 Williams B. A., Rood H. J., 1987, ApJS, 63, 265
 Zabludoff A. I., Mulchaey J. S., 1998, ApJ, 496, 39
 Zabludoff A. I., Mulchaey J. S., 2000, ApJ, 539, 136